Bezari is an extract from turmeric. When sprayed on plants it helps increase resistance to stressful conditions.

Bezari should always be tested on its own on each crop and variety before applying to larger areas to ensure that there is no adverse reaction.

| Composition                 | %    |
|-----------------------------|------|
| Extract of Turmeric Rhizome | 0.5% |
| Potassium Sulphate          | 5%   |
| Adjuvant (carrier)          | Q.S. |
| Total                       | 100% |

#### FIRST AID MEASURES:

Skin contact: Remove contaminated clothing. Wash contaminated area with mild soap & water.

Eye contact: Gently flush eyes with water for at least 15 minutes.

Inhalation: Move person to well ventilated area & seek medical advice immediately.

Ingestion: Seek medical attention if irritation or symptoms persist.

#### LIMITED WARRANTY:

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than that for the purposes indicated on the label. Neither manufacturer nor the seller should be fable for any injury or damage caused by this product as not all posticides and fertilisers have been tested for compatibility. Liability cannot be accepted due to misuse, mishendling or any application not specifically described on the label. Efficacy of any miss will depend upon crop type and growth stage, perticide concerned, climatic conditions, water valuers and various other factors.





Extract of Turmeric Rhizome



FOR AGRICULTURE USE ONLY



# Rate of Application

1 to 2g per litre of water as a foliar application

| сноя                   | RATE/Ha  | TIMING   |
|------------------------|----------|--|
| Vegetables             | 2kg/ha   | After transplantation                                      |
| Horticultural<br>crops | 2-3kg/ha | At basal dose, before flowering, after fruit setting       |
| Pulses                 | 2kg/ha   | At basal dose, before<br>flowering,<br>after ear formation |
| Cereels                | 2kg/ha   | At basal dose, before<br>flowering,<br>after ear formation |
| Ornamental<br>crops    | 1-2kg/ha | At basal dose,<br>before flowering                         |

USE DILUTED SOLUTION IMMEDIATELY AND DO NOT STORE OVERNIGHT.

## HANDLING & STORAGE:

Handling: Avoid contact with eyes and skin. Ensure adequate ventilation of the working area. Use safety gloves, mask, etc.

Storage: Keep out of reach of children, away from food and feed stuffs and animals. Store in a tightly, closed original container and in a dry, dark and well ventilated place.

## Batch Number:

Manufacturing Date:

## **Expiry Date:**

Manufactured for: Earthlee Ventures, UK posta@earthleeventures.com